Neuroscience Letters, 105 (1989) 356–359
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Adler-Graschinsky, E., see Fillinger, E.J., 195
Agid, Y., see Ceballos, I., 41
Aldridge, F., see Lowe, J., 7
Allen, J.M., see Riskind, P.N., 215
Anzai, K., see Horikoshi, T., 340
Asanuma, A., see Horiskohi, T., 340
Audet-Arnold, J., see Riskind, P.N., 215

Beltz, T.G., see Edwards, G.L., 19
Berger, W., see Gollhofer, A., 73
Billy, A.J. and Walters, E.T., Modulation of mechanosensory threshold in *Aplysia* by serotonin, small cardioactive peptide$_B$ (SCP$_B$), FMRFamide, acetylcholine, and dopamine, 200
Bleckmann, H., see Fiebig, E., 57
Blottner, D., Bruggemann, W. and Unsicker, K., Ciliary neurotrophic factor supports target-deprived preganglionic sympathetic spinal cord neurons, 316
Brazell, C., see King, B.H., 174
Brooks, P.A. and Spyer, K.M., Investigation of inositol hexakisphosphate actions in rat nucleus tractus solitarius in vitro, 120
Brtva, R.D., Iwamoto, G.A. and Longhurst, J.C., Distribution of cell bodies for primary afferent fibers from the stomach of the cat, 287
Bruggeman, W., see Blottner, D., 316
Bugiani, O., see Verga, L., 294
Butler, L., see Traynelis, S.F., 326

Ceballos, I., Javoy-Agid, F., Hirsch, E.C., Dumas, S., Kamoun, P.P., Sinet, P.M. and Agid, Y., Localization of copper-zinc superoxide dismutase mRNA in human hippocampus by in situ hybridization, 41
Chabanet, C., see Lavialle, M., 86
Chan, S.H.H., see Chen, C.H., 183
Chan, Y.S., Cheung, Y.M. and Pang, S.F., Elevation of pineal melatonin secretion by electrical stimulation of the cervical sympathetic trunk in rabbits, 107
Chase, T.N., see Mourdian, M.M., 233
Chase, T.N., see Parashos, S.A., 169
Chau, R.M.W., see Jen, L.S., 275
Chen, C.H. and Chan, S.H.H., Involvement of postsynaptic α_2-adrenoceptors and guanine nucleotide-binding protein in guanabenz-induced cardiovascular suppressant effects in the rat, 183
Cheng, G.-F., see Wang, W.-J., 164
Cheung, Y.M., see Chan, Y.S., 107
Chiba, T., Direct synaptic contacts of 5-hydroxytryptamine-, neuropeptide Y-, and somatostatin-immunoreactive nerve terminals on the preganglionic sympathetic neurons of the guinea pig, 281
Ciutat, D., see Esquerda, J.E., 1
Coan, E.J., Irving, A.J. and Collingridge, G.L., Low-frequency activation of the NMDA receptor system can prevent the induction of LTP, 205
Collingridge, G.L., see Coan, E.J., 205
Collins, M.A., see Neafsey, E.J., 344
Comella, J.X., see Esquerda, J.E., 1
Connelly, C.A., Ellenberger, H.H. and Feldman, J.L., Are there serotonergic projections from raphe and retrotrapezoid nuclei to the ventral respiratory group in the rat?, 34
Cunningham, J.T., see Edwards, G.L., 19

Dalm, E., see Gerrits, N.M., 27
Darlington, D.L., Smith, P.F. and Hubbard, J.I., Neuronal activity in the guinea pig medial vestibular nucleus in vitro following chronic unilateral labyrinthectomy, 143
De Arruda Camargo, L.A., see De Luca Junior, L.A., 333
De Bilbao, F., see Senut, M.C., 257
De Luca Junior, L.A., Saad, W.A., De Arruda Camargo, L.A., Renzi, A., Menani, J.V. and Saad, W.A., Carbachol injection into the medial preoptic area induces natriuresis, kaliuresis and antidiuresis in rats, 333
Dederen, P.J.W.C., see Gribnau, A.A.M., 47
Dietz, V., see Gollhofer, A., 73
Dinger, B.G., see Wang, W.-J., 164
Dingledine, R., see Traynelis, S.F., 326
Doherty, F., see Lowe, J., 7
Dourish, C.T., see King, B.H., 174
Drucker, G., see Neafsey, E.J., 344
Dumas, S., see Ceballos, I., 41
Dumortier, B., see Lavialle, M., 86
Dusart, I., Isacson, O., Nothias, F., Gumpel, M.

and Peschanski, M., Presence of Schwann cells in neurodegenerative lesions of the central nervous system, 246

Edwards, G.L., Cunningham, J.T., Beltz, T.G. and Johnson, A.K., Neuropeptide Y-immunoreactive cells in the caudal medulla project to the median preoptic nucleus, 19
Elgoyhen, A.B., see Fillinger, E.J., 195
Ellenberger, H.H., see Connelly, C.A., 34
Epema, A.H., see Gerrits, N.M., 27
Esquerda, J.E., Ciutat, D. and Comella, J.X., Absence of histochemical immunoreactivity to calcitonin gene-related peptide (CGRP) in spinal cord motoneurons from (+)-tubocurarine-treated chick embryos, 1
Evatt, M.L., Wolf, S.L. and Segal, R.L., Modification of human spinal stretch reflexes: preliminary studies, 350

Faille, L., see Martin, A.R., 14
Feldman, J.L., see Connelly, C.A., 34
Ferrell, W.R., see Lam, F.Y., 155
Fidone, S.J., see Wang, W.-J., 164
Fiebig, E. and Bleckmann, H., Cell groups afferent to the telencephalon in a cartilaginous fish (*Platyrhinoidis triseriata*). A WGA-HRP study, 57
Fillinger, E.J., Elgoyen, A.B. and Adler-Graschinsky, E., Benzodiazepines decrease the release of [^{3}H]noradrenaline and of [^{3}H]acetylcholine in the cat superior cervial ganglion, 195
Finnegan, K.T., Skratt, J.J., Irwin, I. and Langston, J.W., The *N*-methyl-D-aspartate (NMDA) receptor antagonist, dextrorphan, prevents the neurotoxic effects of 3,4-methylenedioxymethamphetamine (MDMA) in rats, 300
Frangione, B., see Verga, L., 294

Gabriel, S.M., see Riskind, P.N., 215
Gerrits, N.M., Epema, A.H., Van Linge, A. and Dalm, E., The primary vestibulocerebellar projectin in the rabbit: absence of primary afferents in the flocculus, 27
Giaccone, G., see Vera, L., 294
Gilad, G.M., see Tizabi, Y., 177
Gilad, V.H., see Tizabi, Y., 177
Gill, R. and Kemp, J.A., Protection of CA1 pyramidal cell function by MK-801 following ischaemia in the gerbil, 101
Gollhofer, A., Horstmann, G.A., Berger, W. and Dietz, V., Compensation of translational and rotational perturbations in human posture: stabilization of the centre of gravity, 73
Goto, S., see Horikoshi, T., 340
Grassi, F. and Lux, H.D., Voltage-dependent GABA-induced modulation of calcium currents in chick sensory neurons, 113
Greenberg, D.A., see Jaffe, D.B., 227
Gribnau, A.A.M. and Dederen, P.J.W.C., Collateralization of the cervical corticospinal tract in the rat, 47
Gumpel, M., see Dusart, I., 246

Hamamura, M. and Onaka, T., Pre-weaning handling reduces adrenocorticotropin response to novel but not to noxious stimuli in adult rats, 312
Harrison, N.L. and Lambert, N.A., Modification of $GABA_A$ receptor function by an analog of cyclic AMP, 137
Hatanaka, H., see Hisanaga, K., 321
Hernández, N., see Jaffé, E.H., 189
Heuser, I.J.E., see Mourdian, M.M., 233
Heyes, M.P., see Mourdian, M.M., 233
Hirsch, E.C., see Ceballos, I., 41
Hisanaga, K., Kogure, K., Tsukui, H., Takei, N., Nishio, C. and Hatanaka, H., Increase in choline acetyltransferase activity in septum of rat after transient forebrain ischemia: a possible role of factors released in the hippocampus, 321
Hökfelt, T., see Wiesenfeld-Hallin, Z., 149
Horikoshi, T., Asanuma, A., Yanagisawa, K., Anzai, K. and Goto, S., Regional distribution of metabotropic glutamate response in the rat brain using *Xenopus* oocytes, 340
Horstmann, G.A., see Gollhofer, A., 73
Hubbard, J.I., see Darlington, C.L., 143
Humpel, C. and Saria, A., Effects of GABA and L-glutamic acid on the potassium-evoked in vitro release of substance P- and neurokinin A-like immunoreactivities are different in the rat striatum and substantia nigra, 159

Iino, M., see Tsuzuki, K., 269
Irving, A.J., see Coan, E.J., 205
Irwin, I., see Finnegan, K.T., 300
Isacson, O., see Dusart, I., 246
Ishizuka, K., see Murakamai, T., 96
Iwamoto, G.A., see Brtva, R.D., 287

Jaffe, D.B., Marks, S.S. and Greenberg, D.A., Antagonist drug selectivity for radioligand binding sites on voltage-gated and *N*-methyl-D-aspartate receptor-gated Ca^{2+} channels, 227
Jaffé, E.H. and Hernández, N., Release of [^{3}H]dopamine from rat prefrontal cortex: modulation

through presynaptic cholinergic heteroreceptors, 189
Javoy-Agid, F., see Ceballos, I., 41
Jefferson, D., see Lowe, J., 7
Jen, L.S., Chau, R.M.W. and Tsang, D., Cytochrome oxidase activity in retinas transplanted to the brainstem in rats, 275
Johnson, A.K., see Edwards, G.L., 19

Kalt, G., see Nitsch, C., 263
Kamoun, P.P., see Ceballos, I., 41
Kashima, Y., see Suzuki, T., 211
Kasting, N.W. and Wilinson, M.F., Antipyretic doses of centrally admistered vasopressin reach physiologically meaningful concentrations in the brain of the rat as evaluated by microdialysis, 131
Katayama, N., see Tsuda, M., 241
Kauppila, T., see Pertovaara, A., 68
Kawashima, K., see Suzuki, T., 211
Kemp, J.A., see Gill, R., 101
Kikuchi, K., see Tsuda, M., 241
King, B.H., Brazell, C., Dourish, C.T. and Middlemiss, D.N., MK-212 increases rat plasma ACTH concentration by activation of the 5-HT_{1C} receptor subtype, 174
Kiss, J. and Patel, A.J., Characterization of neurons containing nerve growth factor receptors in the rat neostriatum, 251
Koenig, J.I., see Riskind, P.N., 215
Kogure, K., see Hisanaga, K., 321
Kucera, J. and Walro, J., Alteration of intrafusal bundle composition by nerve crush in neonatal rats, 52

Lam, F.Y. and Ferrell, W.R., Capsaicin suppresses substance P-induced joint inflammation in the rat, 155
Lambert, N.A., see Harrison, N.L., 137
Lamour, Y., see Senut, M.C., 257
Landon, M., see Lowe, J., 7
Langston, J.W., see Finnegan, K.T., 300
Lavialle, M., Chabanet, C. and Dumortier, B., The 24-h rhythm of metabolic activity of the cockroach circadian pacemaker, 86
Lennox, G., see Lowe, J., 7
Longhurst, J.C., see Brtva, R.D., 287
Lothman, E.W., see Stringer, J.L., 91
Lowe, J., Aldridge, F., Lennox, G., Doherty, F., Jefferson, D., Landon, M. and Mayer, R.J., Inclusion bodies in motor cortex and brainstem of patients with motor neurone disease are detected by immunocytochemical localisation of ubiquitin, 7
Lux, H.D., see Grassi, F., 113

MacBride, R.G., see Norvell, J.E., 63
Mallart, A., see Martin, A.R., 14
Marin, C., see Parashos, S.A., 169
Markey, S.P., see Mourdian, M.M., 233
Marks, S.S., see Jaffe, D.B., 227
Martin, A.R., Patel, V., Faille, L. and Mallart, A., Presynaptic calcium currents recorded from calyciform nerve terminals in the lizard ciliary ganglion, 14
Mayer, R.J., see Lowe, J., 7
McNamara, J.O., see Traynelis, S.F., 326
Menani, J.V., see De Luca Junior, L.A., 333
Middlemiss, D.N., see King, B.H., 174
Migheli, A., see Verga, L., 294
Mizutani, N., see Ohya, Y., 221
Morris, R., Responses of spinal dorsal horn neurones evoked by myelinated primary afferent stimulation are blocked by excitatory amino acid antagonists acting at kainate/quisqualate receptors, 79
Mourdian, M.M., Heyes, M.P., Pan, J.-B., Heuser, I.J.E., Markey, S.P. and Chase, T.N., No changes in central quinolinic acid levels in Alzheimer's disease, 233
Murakamai, T., Ishizuka, K. and Uchiyama, M., Convergence of excitatory inputs from the chorda tympani, glossopharyngeal and vagus nerves onto superior salivatory nucleus neurons in the cat, 96

Nagatsu, T., see Ohya, Y., 221
Nakayama, S., see Takaki, M., 125
Naoi, M., see Ohya, Y., 221
Neafsey, E.J., Drucker, G., Raikoff, K. and Collins, M.A., Striatal dopaminergic toxicity following intranigral injection in rats of 2-methylnorharman, a β-carbolinium analog of *N*-methyl-4-phenylpyridinium ion (MPP^+), 344
Nishikawa, T., see Shinohara, K., 307
Nishio, C., see Hisanaga, K., 321
Nitsch, C., Scotti, A., Sommacal, A. and Kalt, G., GABAergic hippocampal neurons resistant to ischemia-induced neornal death contain the Ca^{2+}-binding protein parvalbumin, 263
Norvell, J.E. and MacBride, R.G., Neuropeptide Y (NPY)-like immunoactive nerve fibers in the human and monkey (*Macaca fascicularis*) kidney, 63
Nothias, F., see Dusart, I., 246

Ochi, N., see Ohya, Y., 221

Ohya, Y., Naoi, M., Ochi, N., Mizutani, N., Watanabe, K. and Nagatsu, T., Uptake of *N*-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) and the *N*-methyl-4-phenylpyridinium ion (MPP^+) into fetal mouse brain through the placenta, 221
Onaka, T., see Hamamura, M., 312
Ono, K., see Tsuda, M., 241
Ozawa, S., see Tsuzuki, K., 269

Pan, J.-B., see Mourdian, M.M., 233
Pang, S.F., see Chan, Y.S., 107
Parashos, S.A., Marin, C. and Chase, T.N., Synergy between a selective D_1 antagonist and a selective D_2 antagonist in the induction of catalepsy, 169
Patel, A.J., see Kiss, J., 251
Patel, V., see Martin, A.R., 14
Pertovaara, A. and Kauppila, T., Effect of chronic sciatic nerve section on saphenous nerve input to midline bulboreticular formation in the rat, 68
Peschanski, M., see Dusart, I., 246

Raikoff, K., see Neafsey, E.J., 344
Renzi, A., see De Luca Junior, L.A., 333
Rigsbee, L., see Traynelis, S.F., 326
Riskind, P.N., Allen, J.M., Gabriel, S.M., Koenig, J.I. and Audet-Arnold, J., Sex differences in vasoactive intestinal peptide (VIP) concentrations in the anterior pituitary and hypothalamus of rats, 215

Saad, W.A., see De Luca Junior, L.A., 333
Saria, A., see Humpel, C., 159
Scotti, A., see Nitsch, C., 263
Segal, R.L., see Evatt, M.L., 350
Senut, M.C., De Bilbao, F. and Lamour, Y., Age-related loss of galanin-immunoreactive cells in the rat septal area, 257
Shinohara, K., Nishikawa, T., Yamazaki, K. and Takahashi, K., Ontogeny of strychnine-insensitive [^{3}H]glycine binding sites in rat forebrain, 307
Sinet, P.M., see Ceballos, I., 41
Skratt, J.J., see Finnegan, K.T., 300
Smith, P.F., see Darlington, C.L., 143
Sommacal, A., see Nitsch, C., 263
Spyer, K.M., see Brooks, P.A., 120
Stringer, J.L. and Lothman, E.W., Repetitive seizures cause an increase in paired-pulse inhibition in the dentate gyrus, 91
Suzuki, T., Kashima, Y. and Kawashima, K., Hemicholinium-3-resistant choline uptake system linked to acetylcholine synthesis in the rat hippocampus, 211

Tagliavini, F., see Verga, L., 294
Takahashi, K., see Shinohara, K., 307
Takaki, M. and Nakayama, S., Effects of capsaicin on myenteric neurons of the guinea pig ileum, 125
Takei, N., see Hisanaga, K., 321
Tizabi, Y., Gilad, V.H. and Gilad, G.M., Effects of chronic stressors or corticosterone treatment on the septohippocampal cholinergic system of the rat, 177
Traynelis, S.F., Dingledine, R., McNamara, J.O., Butler, L. and Rigsbee, L., Effect of kindling on potassium-induced electrographic seizures in vitro, 326
Tsang, D., see Jen, L.S., 275
Tsuchiya, T., see Tsuda, M., 241
Tsuda, M., Ono, K., Katayama, N., Yamagata, Y., Kikuchi, K. and Tsuchiya, T., Neurite outgrowth from mouse neuroblastoma and cerebellar cells induced by the protein kinase inhibitor H-7, 241
Tsukui, H., see Hisanaga, K., 321
Tsuzuki, K., Iino, M. and Ozawa, S., Change in calcium permeability caused by quinolinic acid in cultured rat hippocampal neurons, 269

Unsicker, K., see Blottner, D., 316
Uchiyama, M., see Murakamai, T., 96

Van Linge, A., see Gerrits, N.M., 27
Verga, L., Frangione, B., Tagliavini, F., Giaccone, G., Migheli, A. and Bugiani, O., Alzheimer patients and Down patients: cerebral preamyloid deposits differ ultrastructurally and histochemically from the amyloid of senile plaques, 294
Villar, M.J., see Wiesenfeld-Hallin, Z., 149

Walro, J., see Kucera, J., 52
Walters, E.T., see Billy, A.J., 200
Wang, W.-J., Chen, G.-F., Dinger, B.G. and Fidone, S.J., Effects of hypoxia on cyclic nucleotide formation in rabbit carotid body in vitro, 164
Watanabe, K., see Ohya, Y., 221
Wiesenfeld-Hallin, Z., Xu, X.-J., Villar, M.J. and Hökfelt, T., The effect of intrathecal galanin on the flexor reflex in rat: increased depression after sciatic nerve section, 149

Wilkinson, M.F., see Kasting, N.W., 131
Wolf, S.L., see Evatt, M.L., 350

Xu, X.-J., see Wiesenfeld-Hallin, Z., 149

Yamagata, Y., see Tsuda, M., 241
Yamazaki, K., see Shinohara, K., 307
Yanagisawa, K., see Horikoshi, T., 340